

Information about COVID-19 Vaccination

The COVID-19 vaccine you have received is an additional recommended dose to achieve the best possible protection, and for the protection to be as long-lasting as possible.

The vaccine has been assessed to be safe for use. However, just like other vaccines, you may experience some side effects such as headache, body aches, tiredness and soreness at the injection site, or fever. These usually get better after 1-3 days and may be a sign that your immune system is making a protective response against COVID-19.



MINISTRY OF HEALTH SINGAPORE COVID-19 VACCINATION CARD

PERSONAL PARTICULARS

Name: SANDAR
NRIC/FIN: G8840954Q
DOB: 03 Aug 1995

COVID-19 Vaccination Record

S/N	Vaccine Brand / Batch	Date of Vaccination	Place of Vaccination
3	PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / FK8892	09 Jan 2022	Raffles Medical VC - Sengkang CC

Past COVID-19 Vaccination Record(s)

- 11 Jul 2021 / PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / 1E022A / RAFFLES MEDICAL VACCINATION CENTRE @ SENGKANG CC
- 14 Aug 2021 / PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / FF2154 / RAFFLES MEDICAL VACCINATION CENTRE @ SENGKANG CC

After Vaccination Advice

Possible Side Effects	How to Manage
Pain, redness, swelling at the injection site	Those with fever are advised to self-isolate at home until the fever subsides.
Fever, chills	Paracetamol 1 to 2 tablets every 6 hours for adults or dosed according to the child's weight as needed.
Headache, muscle pain, joint pain	Rest.
Tiredness	Usually gets better by itself in a week or so.
Lymph node swelling at neck or arms	

See a doctor if:

- the side effects persist or get worse
 - the fever persists for more than 48 hours
- You can also report any side effects that you experience to HSA via go.gov.sg/hf
- This vaccine may cause a severe allergic reaction in very rare instances. If you experience a severe allergic reaction (difficulty breathing, swelling of your face, throat, eyes or lips, a fast heartbeat, dizziness or weakness, rash all over your body), please seek immediate medical attention by calling 995 or going to the nearest A&E. The doctor will also decide if you should receive the second dose of the vaccine.
- If you experience chest pain, difficulty breathing or abnormal heartbeats, please also seek medical attention immediately.

Information about COVID-19 Vaccination

The COVID-19 vaccine you have received consists of two doses. The second dose must be completed to achieve the best possible protection, and for the protection to be as long-lasting as possible.

The vaccine has been assessed to be safe for use. However, just like other vaccines, you may experience some side effects such as headache, body aches, tiredness and soreness at the injection site, or fever. These usually get better after 1-3 days and may be a sign that your immune system is making a protective response against COVID-19.



MINISTRY OF HEALTH SINGAPORE COVID-19 VACCINATION CARD

Please bring this card for your next appointment

PERSONAL PARTICULARS

Name: SANDAR
NRIC/FIN: G8840954Q
DOB: 03 Aug 1995

Your Next Vaccination is Due in _____ Days

Appointment Date	Time	Place
14 Aug 2021	01:00 PM	Raffles Medical VC - Sengkang CC

COVID-19 Vaccination Record

Vaccine Dose	Vaccine Brand / Batch	Date of Vaccination	Place of Vaccination
D1	PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / 1E022A	11 Jul 2021	Raffles Medical VC - Sengkang CC

After Vaccination Advice

Possible Side Effects	How to Manage
Pain, redness, swelling at the injection site	Those with fever are advised to self-isolate at home until the fever subsides.
Fever, chills	Paracetamol 1 to 2 tablets every 6 hours for adults or dosed according to the child's weight as needed.
Headache, muscle pain, joint pain	Rest.
Tiredness	Usually gets better by itself in a week or so.
Lymph node swelling at neck or arms	

See a doctor if:

- the side effects persist or get worse
 - the fever persists for more than 48 hours
- You can also report any side effects that you experience to HSA via go.gov.sg/hf
- This vaccine may cause a severe allergic reaction in very rare instances. If you experience a severe allergic reaction (difficulty breathing, swelling of your face, throat, eyes or lips, a fast heartbeat, dizziness or weakness, rash all over your body), please seek immediate medical attention by calling 995 or going to the nearest A&E. The doctor will also decide if you should receive the second dose of the vaccine.

Information about COVID-19 Vaccination

The COVID-19 vaccine you have received consists of two doses. The second dose must be completed to achieve the best possible protection, and for the protection to be as long-lasting as possible.

The vaccine has been assessed to be safe for use. However, just like other vaccines, you may experience some side effects such as headache, body aches, tiredness and soreness at the injection site, or fever. These usually get better after 1-3 days and may be a sign that your immune system is making a protective response against COVID-19.



MINISTRY OF HEALTH SINGAPORE COVID-19 VACCINATION CARD

Please bring this card for your next appointment

PERSONAL PARTICULARS

Name: SANDAR
NRIC/FIN: G8840954Q
DOB: 03 Aug 1995

Your Next Vaccination is Due in _____ Days

Appointment Date	Time	Place

COVID-19 Vaccination Record

Vaccine Dose	Vaccine Brand / Batch	Date of Vaccination	Place of Vaccination
D1	PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / 1E022A	11 Jul 2021	Raffles Medical VC - Sengkang CC
D2	PFIZER-BIONTECH/COMIRNATY COVID-19 Vaccine / FF2154	14 Aug 2021	Raffles Medical VC - Sengkang CC

After Vaccination Advice

Possible Side Effects	How to Manage
Pain, redness, swelling at the injection site	Those with fever are advised to self-isolate at home until the fever subsides.
Fever, chills	Paracetamol 1 to 2 tablets every 6 hours for adults or dosed according to the child's weight as needed.
Headache, muscle pain, joint pain	Rest.
Tiredness	Usually gets better by itself in a week or so.
Lymph node swelling at neck or arms	

See a doctor if:

- the side effects persist or get worse
 - the fever persists for more than 48 hours
- You can also report any side effects that you experience to HSA via go.gov.sg/hf
- This vaccine may cause a severe allergic reaction in very rare instances. If you experience a severe allergic reaction (difficulty breathing, swelling of your face, throat, eyes or lips, a fast heartbeat, dizziness or weakness, rash all over your body), please seek immediate medical attention by calling 995 or going to the nearest A&E. The doctor will also decide if you should receive the second dose of the vaccine.
- If you experience chest pain, difficulty breathing or abnormal heartbeats, please also seek medical attention immediately.